


AMENDMENTS TO THE CLAIMS

This listing of claims replaces all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS


1. (original) A 1,3,5-triazine compound represented by the following formula I:



wherein R^1 and R^2 are each independently a methyl group, an ethyl group, a hydroxyalkyl group having 2 to 5 carbon atoms, $-(CH_2CH_2O)_mR^6$ (where m is an integer of 1 to 120, and R^6 is a hydrogen atom, a methyl group, an ethyl group, or a propyl group), $-(CH_2CH_2NR^7)_mH$ (where m is an integer of 1 to 120, and R^7 is an alkyl group having 2 to 5 carbon atoms, an N,N-dialkylaminoethyl group or $-CH_2CH_2N^+(CH_3)_3$), $-CH_2CH_2SO_3^-$, $-CH_2CH_2N^+(CH_3)_3$, or an alkyl group having 6 to 20 carbon atoms, but both R^1 and R^2 are not alkyl groups having 6 to 20 carbon atoms at the same time; one or two of R^3 , R^4 and R^5 are methyl groups, and the remaining R^3 , R^4 and R^5 are each independently $-CH_2COO-C_nH_{2n+1}$, $-C_nH_{2n+1}$, or $-C_6H_4-p-C_nH_{2n+1}$, where n is an integer of 6 to 20, and $-C_nH_{2n+1}$ is linear; and X^- is a halide ion, a triflate anion, a nitrate ion, a sulfate ion, a hydrogensulfate ion, a sulfonate ion, a tetrafluoroborate ion, or a perchlorate ion.


2. (original) The compound of claim 1, wherein at least one of R^1 and R^2 is a methyl group or an ethyl group.

3. (currently amended) The compound of claim 1 or 2, wherein n is 12 to 16.
4. (original) A method for producing a 1,3,5-triazine compound represented by the following formula I':




wherein R^1 and R^2 are each independently a methyl group, an ethyl group, a hydroxyalkyl group having 2 to 5 carbon atoms, $-(CH_2CH_2O)_mR^6$ (where m is an integer of 1 to 120, and R^6 is a hydrogen atom, a methyl group, an ethyl group, or a propyl group), $-(CH_2CH_2NR^7)_mH$ (where m is an integer of 1 to 120, and R^7 is an alkyl group having 2 to 5 carbon atoms, an N,N-dialkylaminoethyl group or $-CH_2CH_2N^+(CH_3)_3$, $-CH_2CH_2SO_3^-$, $-CH_2CH_2N^+(CH_3)_3$, or an alkyl group having 6 to 20 carbon atoms, but both R^1 and R^2 are not alkyl groups having 6 to 20 carbon atoms at the same time; one or two of R^3 , R^4 and R^5 are methyl groups, and the remaining R^3 , R^4 and R^5 are each independently $-CH_2COO-C_nH_{2n+1}$, $-C_nH_{2n+1}$, or $-C_6H_4-p-C_nH_{2n+1}$, where n is an integer of 6 to 20, and $-C_nH_{2n+1}$ is linear; and X^- is a triflate anion, comprising:

obtaining triflate by mixing a compound represented by the following formula II and trifluoromethanesulfonic anhydride in an organic solvent:




wherein R^1 and R^2 are each independently a methyl group, an ethyl group, a hydroxyalkyl group having 2 to 5 carbon atoms, $-(CH_2CH_2O)_mR^6$ (where m is an integer of 1 to 120, and R^6 is a hydrogen atom, a methyl group, an ethyl group, or a propyl group), $-(CH_2CH_2NR^7)_mH$ (where m is an integer of 1 to 120, and R^7 is an alkyl group having 2 to 5 carbon atoms, an N,N-dialkylaminoethyl group or $-CH_2CH_2N^+(CH_3)_3$), $-CH_2CH_2SO_3^-$, $-CH_2CH_2N^+(CH_3)_3$, or an alkyl group having 6 to 20 carbon atoms, but both R^1 and R^2 are not alkyl groups having 6 to 20 carbon atoms at the same time; and mixing the obtained triflate and a tertiary amine represented by the following formula III in an appropriate organic solvent:



wherein one or two of R^3 , R^4 and R^5 are methyl groups, and the remaining R^3 , R^4 and R^5 are each independently $-CH_2COO-C_nH_{2n+1}$, $-C_nH_{2n+1}$, or $-C_6H_4-p-C_nH_{2n+1}$, where n is an integer of 6 to 20, and $-C_nH_{2n+1}$ is linear.


5. (original) A method for producing a 1,3,5-triazine compound represented by the following formula I'':




wherein R^1 and R^2 are each independently a methyl group, an ethyl group, a hydroxyalkyl group having 2 to 5 carbon atoms, $-(CH_2CH_2O)_mR^6$ (where m is an integer of 1 to 120, and R^6 is a hydrogen atom, a methyl group, an ethyl group, or a propyl

group), $-(\text{CH}_2\text{CH}_2\text{NR}^7)_m\text{H}$ (where m is an integer of 1 to 120, and R^7 is an alkyl group having 2 to 5 carbon atoms, an N,N-dialkylaminoethyl group or $-\text{CH}_2\text{CH}_2\text{N}^+(\text{CH}_3)_3$, $-\text{CH}_2\text{CH}_2\text{SO}_3^-$, $-\text{CH}_2\text{CH}_2\text{N}^+(\text{CH}_3)_3$, or an alkyl group having 6 to 20 carbon atoms, but both R^1 and R^2 are not alkyl groups having 6 to 20 carbon atoms at the same time; one or two of R^3 , R^4 and R^5 are methyl groups, and the remaining R^3 , R^4 and R^5 are each independently $-\text{CH}_2\text{COO}-\text{C}_n\text{H}_{2n+1}$, $-\text{C}_n\text{H}_{2n+1}$, or $-\text{C}_6\text{H}_4\text{-p-C}_n\text{H}_{2n+1}$, where n is an integer of 6 to 20, and $-\text{C}_n\text{H}_{2n+1}$ is linear; and X^- is a halide ion, comprising:

mixing a compound represented by the following formula IV and a tertiary amine represented by the following formula III in an appropriate solvent:



wherein R^1 and R^2 are each independently a methyl group, an ethyl group, a hydroxyalkyl group having 2 to 5 carbon atoms, $-(\text{CH}_2\text{CH}_2\text{O})_m\text{R}^6$ (where m is an integer of 1 to 120, and R^6 is a hydrogen atom, a methyl group, an ethyl group, or a propyl group), $-(\text{CH}_2\text{CH}_2\text{NR}^7)_m\text{H}$ (where m is an integer of 1 to 120, and R^7 is an alkyl group having 2 to 5 carbon atoms, an N,N-dialkylaminoethyl group or $-\text{CH}_2\text{CH}_2\text{N}^+(\text{CH}_3)_3$, $-\text{CH}_2\text{CH}_2\text{SO}_3^-$, $-\text{CH}_2\text{CH}_2\text{N}^+(\text{CH}_3)_3$, or an alkyl group having 6 to 20 carbon atoms, but both R^1 and R^2 are not alkyl groups having 6 to 20 carbon atoms at the same time; and X is a halogen atom;




wherein one or two of R^3 , R^4 and R^5 are methyl groups, and the remaining R^3 , R^4

and R^5 are each independently $-\text{CH}_2\text{COO}-\text{C}_n\text{H}_{2n+1}$, $-\text{C}_n\text{H}_{2n+1}$, or $-\text{C}_6\text{H}_4\text{-p-C}_n\text{H}_{2n+1}$, where n is an integer of 6 to 20, and $-\text{C}_n\text{H}_{2n+1}$ is linear.

6. (original) A method for producing a carboxylic acid derivative, comprising:


mixing a carboxylic acid and a compound having a nucleophilic functional group in an aqueous solution in the presence of a 1,3,5-triazine compound represented by the following formula I:



wherein R^1 and R^2 are each independently a methyl group, an ethyl group, a hydroxyalkyl group having 2 to 5 carbon atoms, $-(\text{CH}_2\text{CH}_2\text{O})_m\text{R}^6$ (where m is an integer of 1 to 120, and R^6 is a hydrogen atom, a methyl group, an ethyl group, or a propyl group), $-(\text{CH}_2\text{CH}_2\text{NR}^7)_m\text{H}$ (where m is an integer of 1 to 120, and R^7 is an alkyl group having 2 to 5 carbon atoms, an N,N-dialkylaminoethyl group or $-\text{CH}_2\text{CH}_2\text{N}^+(\text{CH}_3)_3$), $-\text{CH}_2\text{CH}_2\text{SO}_3^-$, $-\text{CH}_2\text{CH}_2\text{N}^+(\text{CH}_3)_3$, or an alkyl group having 6 to 20 carbon atoms, but both R^1 and R^2 are not alkyl groups having 6 to 20 carbon atoms at the same time; one or two of R^3 , R^4 and R^5 are methyl groups, and the remaining R^3 , R^4 and R^5 are each independently $-\text{CH}_2\text{COO}-\text{C}_n\text{H}_{2n+1}$, $-\text{C}_n\text{H}_{2n+1}$, or $-\text{C}_6\text{H}_4\text{-p-C}_n\text{H}_{2n+1}$, where n is an integer of 6 to 20, and $-\text{C}_n\text{H}_{2n+1}$ is linear; and X^- is a halide ion, a triflate anion, a nitrate ion, a sulfate ion, a hydrogensulfate ion, a sulfonate ion, a tetrafluoroborate ion, or a perchlorate ion.

7. (original) The method of claim 6, wherein the carboxylic acid is a fatty acid having 6 to 20 carbon atoms.


8. (original) The method of claim 7, wherein the carboxylic acid is a fatty acid having 8 to 18 carbon atoms.
9. (currently amended) The method of claim 6 ~~any of claims 6 to 8~~, wherein at least one of R^1 and R^2 in the formula I is a methyl group or an ethyl group.
10. (currently amended) The method of claim 6 ~~any of claims 6 to 9~~, wherein n in the formula I is 12 to 16.
11. (currently amended) The method of claim 6 ~~any of claims 6 to 10~~, wherein the compound having a nucleophilic functional group is a primary amine compound or a secondary amine compound.
12. (original) A method for producing a carboxylic acid derivative, comprising mixing:
 a carboxylic acid;
 a compound having a nucleophilic functional group;
 a compound represented by the following formula IV; and
 a tertiary amine represented by the following formula III in an aqueous solution:



(IV)

wherein R^1 and R^2 are each independently a methyl group, an ethyl group, a hydroxyalkyl group having 2 to 5 carbon atoms, $-(CH_2CH_2O)_mR^6$ (where m is an integer of 1 to 120, and R^6 is a hydrogen atom, a methyl group, an ethyl group, or a propyl group), $-(CH_2CH_2NR^7)_mH$ (where m is an integer of 1 to 120, and R^7 is an alkyl group having 2 to 5 carbon atoms, an N,N-dialkylaminoethyl group or $-CH_2CH_2N^+(CH_3)_3$), -

$\text{CH}_2\text{CH}_2\text{SO}_3^-$, $-\text{CH}_2\text{CH}_2\text{N}^+(\text{CH}_3)_3$, or an alkyl group having 6 to 20 carbon atoms, but both R^1 and R^2 are not alkyl groups having 6 to 20 carbon atoms at the same time; and X is a halogen atom,



wherein one or two of R^3 , R^4 and R^5 are methyl groups, and the remaining R^3 , R^4 and R^5 are each independently $-\text{CH}_2\text{COO}-\text{C}_n\text{H}_{2n+1}$, $-\text{C}_n\text{H}_{2n+1}$, or $-\text{C}_6\text{H}_4\text{-p}-\text{C}_n\text{H}_{2n+1}$, where n is an integer of 6 to 20, and $-\text{C}_n\text{H}_{2n+1}$ is linear.

13. (original) The method of claim 12, wherein the carboxylic acid is a fatty acid having 6 to 20 carbon atoms.

14. (original) The method of claim 13, wherein the carboxylic acid is a fatty acid having 8 to 18 carbon atoms.

15. (currently amended) The method of claim 12 ~~any of claims 12 to 14~~, wherein at least one of R^1 and R^2 in the formula I is a methyl group or an ethyl group.

16. (currently amended) The method of claim 12 ~~any of claims 12 to 15~~, wherein n in the formula III is 12 to 16.

17. (currently amended) The method of claim 12 ~~any of claims 12 to 16~~, wherein the compound having a nucleophilic functional group is a primary amine compound or secondary amine compound.

18. (currently amended) The method of claim 12 ~~any of claims 12 to 16~~, wherein the compound having a nucleophilic functional group is an alcohol compound.